

L18 ANSWER 7 OF 8

MEDLINE

ACCESSION NUMBER: 95053743 MEDLINE
DOCUMENT NUMBER: 95053743 PubMed ID: 7964486
TITLE: Mechanisms of **tolerance** induction in major histocompatibility complex class II-restricted T cells specific for a blood-borne self-antigen.
AUTHOR: Zal T; Volkmann A; Stockinger B
CORPORATE SOURCE: Department of Molecular Immunology, National Institute for Medical Research, Mill Hill, London, United Kingdom.
SOURCE: JOURNAL OF EXPERIMENTAL MEDICINE, (1994 Dec 1) 180 (6) 2089-99.
Journal code: 2985109R. ISSN: 0022-1007.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals; AIDS
ENTRY MONTH: 199412
ENTRY DATE: Entered STN: 19950110
Last Updated on STN: 19950110
Entered Medline: 19941223

AB Transgenic mice expressing a major histocompatibility complex class II-restricted T cell receptor with specificity for a natural self-antigen,

the fifth component of complement, were generated to analyze the mechanism

of **tolerance** induction to a blood-borne self-protein. In the absence of C5 protein thymocytes from T cell receptor transgenic mice develop into mature **CD4** single positive cells which emigrate into the periphery and mount C5-specific T cell responses upon immunization with C5. In the presence of circulating C5 protein, **CD4** single positive thymocytes do not develop. Negative selection occurs late in thymic ontogeny leaving the bulk of **CD4+8+** thymocytes unaffected. This phenotype may be due to a delay in contact with self-antigen presentation which, under physiological conditions, is inefficient in the cortex of C5+ mice, and therefore does not affect most immature double positive thymocytes. In contrast, in vitro exposure to C5(-)-presenting **dendritic** cells or in vivo injection of C5 peptide results in deletion of double positive thymocytes. C5+ transgenic mice are tolerant in vivo, but contain T cells in spleen and lymph nodes that secrete interleukin 2 and interferon gamma in response to C5 activation in vitro. When crossed onto a Rag1-/- background to prevent endogenous T cell receptor rearrangements, these peripheral potentially autoreactive cells do not appear. This indicates that endogenous T cell receptor rearrangements possibly leading to the expression of two receptors might be a prerequisite for their survival and export into the periphery.